

Xencongridae

by J. Blache, M.-L. Bauchot & L. Saldanha

G: 1
Sp: 1

- 77.1 **CHLOPSIS** Rafinesque, 1810 Gender: F
Indice d'Ittiologia siciliana . . . : 42, 58 (type: *C. bicolor* Rafinesque, 1810, by monotypy).

- 77.1.1 **Chlopsis bicolor** Rafinesque, 1810
Chlopsis bicolor Rafinesque, 1810b, Indice d'Ittiologia siciliana . . . : 58, pl. 2 (fig. 1) ('Sicilia'). Type: unknown.
Chlopsis bicolor: Carus, 1889–1893 : 545 | Grassi & Calandruccio, 1894b : 7; 1896b : 349; 1897f : 43 | Facciola, 1901a : 59 | Supino, 1905a : 25, fig. 1–9 | Grassi, 1913 : 130, pl. 7 (fig. 31–39) | Facciola, 1914 : 39 | Kirincic & Lepetic, 1953 : 1–2 | Parenzan, 1956 : 297 | Šoljan, 1963 : 395 | Tortonese, 1963 : 165; 1970e : 316, fig. 125–128.
Eggs, larvae and young stages. Strömman, 1896 : 39, pl. 5 (fig. 5–6) | Sanzo, 1905 : 315 | Schmidt, 1912b : 338; 1912e : 53, 1 fig. | Grassi, 1912 : 17, 1 fig.; 1913 : 130, pl. 7 (fig. 31–39), pl. 10 (fig. 22), pl. 14 (fig. 3–5); 1915 : 702 | Bertin, 1926: tab. 6 | Fish, 1927 : 307 | Ancona, 1931 : 115, fig. 86–91 | Sparta, 1939a : 3, fig. 1–10 | Orton, 1964 : 438 | Smith, 1969 : 390.
Otoliths (sagitta). No data.
Habitat, distribution and abundance. Benthic on the continental slope. Mediterranean, Eastern Atlantic south to Angola. Not very common.
Common names. None.