

Nettodaridae

by J. Blache, M.-L. Bauchot & L. Saldanha

G: 1
Sp: 1

- 81.1 **NETTODARUS** Whitley, 1951 Gender: M
Rec. Aust. Mus., **22** : 407 (type: *Nettastoma brevirostre*, Facciola, 1887, by orig. design.).
Todarus Grassi & Calandruccio, 1896, *Atti Accad. naz. Lincei Rc.*, **5** : 349, preoccupied.
- 81.1.1 *Nettodarus brevirostris* (Facciola, 1887)
Nettastoma brevirostre Facciola, 1887, *Naturalista sicil.*, **6** (9) : 166, pl. 3 (fig. 3) ('Sicilia'). Type lost.
Nettastoma brevirostre: Carus, 1893 : 543.
Nettastoma brevirostris: Griffini, 1903 : 171.
Todarus brevirostris: Supino, 1905c : 255 | Grassi, 1913 : 170, pl. 10 (fig. 11–17) | Trotti, 1947 : 1; 1948 : 93 | Tortonese, 1963 : 165.
Nettodarus brevirostris: Tortonese, 1970 : 337, fig. 136 | Blache *et al.*, 1970b : 319–331, 15 fig.
 Eggs, larvae and young stages. Grassi, 1912 : 18; 1913 : 170, pl. 10 (fig. 1–17) | Schmidt, 1913 : 12, pl. 1 (fig. 8) (*Leptocephalus telescopicus*) | Grassi, 1917 : 25, pl. 1 (fig. 10–12) | Ancona, 1931 : 123, fig. 102–103 | Castle, 1969 : 24, 55.
 Otoliths (sagitta). No data.
 Habitat, distribution and abundance. Benthic, on the continental slope. Mediterranean, western and eastern Atlantic (south to 5° S.). Rare.