

85

# Nessorhamphidae

by J. Blache, M.-L. Bauchot & L. Saldanha

G: 1

Sp: 1

- 85.1 **NESSORHAMPHUS** Schmidt, 1930 Gender: M  
*Vidensk. Meddr dansk naturh. Foren., Kbh.*, **90** : 372 (type: *Leptocephalus ingolfianus* Schmidt, 1912, by monotypy).
- 85.1.1 *Nessorhamphus ingolfianus* (Schmidt, 1912)  
*Leptocephalus ingolfianus* Schmidt, 1912d, *Vidensk. Meddr dansk naturh. Foren., Kbh.*, **64** : 49, pl. 3 (fig. 8) ('S.W. of the Azores: 32° 03' N., 39 00' W.), larval stage. Syn-  
types: ZMUC no. P 32520–23.  
*Nessorhamphus ingolfianus*: Schmidt, 1930, *Vidensk. Meddr dansk naturh. Foren., Kbh.*, **90** : 373 ('warmer parts of all oceans').  
*Nessorhamphus ingolfianus*: Schmidt, 1931b : 487; 1932b : 251, fig. 192.  
Eggs, larvae and young stages. Schmidt, 1912b : 340; 1912d : 49, pl. 3 (fig. 8) | Ancona, 1928 : 518 | Schmidt, 1930 : 373, pl. 5; 1931a : 444, fig. 3 | Beebe, 1935 : 25, 29, fig. 10 | Bruun, 1936 : 10, fig. 4.  
Otoliths (sagitta). No data.  
Habitat, distribution and abundance. Epi- to mesopelagic. Cosmopolite. Common.  
Common names. None.